

Black Acara (*Cichlasoma bimaculatum*)

Ecological Risk Screening Summary

Web Version – 10/18/2012



Photo: F. Naneix

1 Native Range and Nonindigenous Occurrences

Native Range

From Nico et al. (2012):

“Tropical America. South America in the Guianas from Moruka River in Guyana east to about Sinnamary in French Guiana; also in Essequibo basin, and in adjacent Rio Branco drainage of Amazon basin (Kullander and Nijssen 1989).”

Nonindigenous Occurrences

From Nico et al. (2012):

“The species has been established in Florida, since the early 1960s; it was first discovered in Dade County (Rivas 1965). More recently it was documented at Flint Penn Strand (Ceilley and Ceilley, 2000). The expanded geographic range of the species includes the counties of:”

- “Broward (Courtenay et al. 1974; Courtenay and Hensley 1979a);
- Collier (Courtenay and Hensley 1979a; Courtenay et al. 1986);
- Dade (Kushlan 1972; Courtenay et al. 1974; Hogg 1976; Courtenay and Hensley 1979a; Loftus and Kushlan 1987);
- Hendry (Courtenay and Hensley 1979a);
- Lee (Nico, unpublished data);
- Martin (museum specimens);
- Monroe (Kushlan 1972; Courtenay et al. 1974; Courtenay and Hensley 1979a; Loftus and Kushlan 1987), and
- Palm Beach (Courtenay et al. 1974; Courtenay and Hensley 1979a).”

“There are unconfirmed reports or possible occurrences of this species in Glades and Okeechobee counties (Courtenay et al. 1986; Courtenay and Stauffer 1990). It is established in Big Cypress National Preserve and in Everglades National Park (Kushlan 1972; Loftus and Kushlan 1987; Lorenz et al. 1997; Tilmant 1999; Lovtus 2004). It is also established in Florida Panther National Wildlife Refuge (USFWS 2005).”

Means of Introductions

From Nico et al. (2012):

“In southeastern Florida, introduction was the result of escapes and intentional releases from fish farms, probably during the mid- to late 1950s (Courtenay and Hensley 1979a). Some releases were deliberate attempts to dispose of unwanted and sometimes illegal fish stocks (Courtenay and Stauffer 1990).”

Remarks

From Nico et al. (2012): “Established in Florida.”

“The first Florida specimens of *C. bimaculatum* were reported and identified as *Aequidens portalegrensis* (= *Cichlasoma portalegreense*) by Rivas (1965), Bailey et al. (1970), Kushlan (1972), and others. Cultured in Florida possibly since the 1930s, *Cichlasoma bimaculatum* remained an important part of the aquarium trade until the late 1950s, and probably was the first aquarium fish to become established in open waters of Florida (Courtenay and Stauffer 1990). One of the most abundant introduced species in south Florida canal systems during the early 1970s, *C. bimaculatum* has been gradually replaced by *Tilapia mariae* as the most abundant cichlid, possibly through competition for space (Courtenay and Hensley 1979a, 1979b; Loftus and Kushlan 1987). Loftus and Kushlan (1987) provided a map showing the species' distribution in south Florida. Throughout its south Florida range, it is much more common in disturbed habitats, mainly canals, than in natural habitats (Loftus and Kushlan 1987). Spawning has been observed during every month of the year in southeastern Florida (Courtenay 1989).”

2 Biology and Ecology

Taxonomic Hierarchy

From ITIS (2012):

Kingdom Animalia
 Phylum Chordata
 Subphylum Vertebrata
 Superclass Osteichthyes
 Class Actinopterygii
 Subclass Neopterygii
 Infraclass Teleostei
 Superorder Acanthopterygii
 Order Perciformes
 Suborder Labroidei
 Family Cichlidae
 Genus *Cichlasoma*
 Species *Cichlasoma bimaculatum*

Taxonomic standing: valid

Size, Weight, Age

From Froese and Pauly (2010):

“Max length : 12.3 cm SL male/unsexed; (Kullander 2003); 30 cm TL (female); common length : 12.0 cm TL male/unsexed; (Hugg 1996)”

Environment

From Froese and Pauly (2010):

Benthopelagic; Freshwater (catadromous); pH range: 6.5 – 7.0

Climate/Range

From Froese and Pauly (2010): Tropical; 16°C - 24°C (Riehl and Baensch 1996); 12°N - 1°N

Distribution

From Froese and Pauly (2010):

“South America: Orinoco River basin, in the Caroni in River Venezuela; Guianas, from the Essequibo River to the Sinnamary River; Amazon River basin, in the upper Branco River basin.”

Biology

From Froese and Pauly (2010):

“Occurs in canals and swamps (Mills and Vevers 1989). Tolerates low oxygen. Feeds on crustaceans and insects (Mills and Vevers 1989). Males are bigger than females which become mature at 7 to 9 cm (20-25 g). Spawning occurs at the start of the rainy season. About 700 eggs are spawned on open substrate and are cared for by the parents. Max length 35 cm TL (Ref. 35237).”

Human uses

From Froese and Pauly (2010):

“Fisheries: of no interest; gamefish: yes; aquarium: commercial”

Diseases

From Froese and Pauly (2010):

“*Mycobacterium chelonae abscessus*”

“*Mycobacterium fortuitum*”

“*Mycobacterium simiae*”

C. bimaculatum is known to harbor and transmit bacterial pathogens that can cause a fatal disease (tuberculosis) among a wide variety of fish, as well as irritating skin lesions and pulmonary complications in humans (Gray et al. 1990, Howard et al. 1987, Landsell et al. 1993, Nigrelli et al. 1963, and Parisot et al. 1970).

Threat to humans

From Froese and Pauly (2010): “Disease vector”

3 Impacts of Introductions

Considered a competitor with native sunfish for spawning areas; juvenile bands of *C. bimaculatum* have been observed to surround spawning bluegill *Lepomis macrochirus* aggressively in attempts to feed on eggs (Hogg 1976).

4 Global Distribution




Figure 1 (above). Global distribution of *C. bimaculatum*. Map from GBIF (2010). The location in Oregon (United States) was not used as it does not represent an established population.

5 Distribution within the United States



Figure 2 (above). Distribution of *C. bimaculatum* in the United States. Map from Nico et al. (2012).

6 CLIMATCH

Summary of Climate Matching Analysis

The climate match (Australian Bureau of Rural Science 2010; 16 climate variables; Euclidean Distance) was high in Florida, some of the Great Lakes States, and along the West Coast. Medium matches covered the rest of the country except for the Northern Plains States. Climate 6 match indicated that the US has a high climate match. The range for a high climate match is 0.103 and greater; the climate match of *C. bimaculatum* is 0.223.




Figure 3 (above). CLIMATCH (Australian Bureau of Rural Science 2010) source map showing weather stations selected as source locations (red) and non-source locations (blue) for *C. bimaculatum* climate matching. Source locations from GBIF (2010) and Nico et al. (2012).




Figure 4 (above). Map of CLIMATCH (Australian Bureau of Rural Science 2010) climate matches for *C. bimaculatum* in the continental United States based on source locations reported by GBIF (2010) and Nico et al. (2012).. 0= Lowest match, 10=Highest match.

Table 1 (below). CLIMATCH (Australian Bureau of Rural Science 2010) climate match scores.

CLIMATCH Score	0	1	2	3	4	5	6	7	8	9	10
Count	1	12	65	221	629	605	295	63	40	5	38
Climate 6 Proportion =			0.223	(High)							

7 Certainty of Assessment

Information on the biology of *C. bimaculatum* is fairly abundant. However, there is very little information available on the impacts caused by introduction of this species. In order for higher certainty more research is needed. Certainty of this assessment is medium.

8 Risk Assessment

Summary of Risk to the Continental United States

C. bimaculatum is established in Florida, spawns throughout the year, and competes for spawning habitat with native species of fish in south Florida drainage canals at trust resource sites including national parks, national preserves, and national wildlife refuges. *C. bimaculatum* is known to harbor and transmit bacterial pathogens that can cause a fatal disease (tuberculosis) among a wide variety of fish, as well as irritating skin lesions and pulmonary complications in humans (Gray et al. 1990, Howard et al. 1987, Landsell et al. 1993, Nigrelli et al. 1963, and Parisot et al. 1970). The overall risk of this species is considered high due its disease carrying aspects (See Sec. 2 and 8).

Assessment Elements

- **History of Invasiveness (Sec. 3):** High
- **Climate Match (Sec. 6):** High
- **Certainty of Assessment (Sec. 7):** Medium
- **Remarks:** Potential disease vector to native fish and humans.
- **Overall Risk Assessment Category:** High

9 References

Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.

Australian Bureau of Rural Sciences. 2010. CLIMATCH. Available:
<http://adl.brs.gov.au:8080/Climatch/> (Accessed August 3, 2010).

Froese, R. and D. Pauly (Eds). 2010. *Cichlasoma bimaculatum*. FishBase (www.fishbase.org, version 03/2010). Available:
<http://www.fishbase.org/Summary/speciesSummary.php?ID=3613&genusname=Cichlasoma&speciesname=bimaculatum&lang=English> (Accessed 2 August 2010).

GBIF. 2010. *Cichlasoma bimaculatum*. Global Biodiversity Information Facility. Available:
<http://data.gbif.org/species/13578228/> (Accessed August 2, 2010).

ITIS. 2012. *Cichlasoma bimaculatum*. Integrated Taxonomic Information System Online Database (<http://www.itis.gov/>). Available:
http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=169775 (Accessed October 10, 2012).

Nico, L., P. Fuller, and M. Neilson. 2012. *Cichlasoma bimaculatum*. USGS Nonindigenous Aquatic Species Database, Gainesville, FL. Available:
<http://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=441> (Accessed October 10, 2012)

10 References Quoted But Not Accessed

Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.

Bailey, R. M., J. E. Fitch, E. S. Herald, E. A. Lachner, C. C. Lindsey, C. R. Robins, and W. B. Scott. 1970. A list of common and scientific names of fishes from the United States and Canada, 3rd edition. American Fisheries Society Special Publication 6. American Fisheries Society, Bethesda, MD.

Ceilley, D.W. and D.E. Ceilley. 2000. A survey of freshwater fishes in the hydric flatwoods of Flint Pen Strand, Lee County, Florida. unpublished manuscript.

Courtenay, W. R., Jr. 1989. Exotic fishes in the National Park System. Pages 237-252 in L. K. Thomas, editor. Proceedings of the 1986 conference on science in the national parks, volume 5. Management of exotic species in natural communities. U.S. National Park Service and George Wright Society, Washington, DC.

Courtenay, W. R., Jr., and D. A. Hensley. 1979a. Survey of introduced non-native fishes. Phase I Report. Introduced exotic fishes in North America: status 1979. Report Submitted to National Fishery Research Laboratory, U.S. Fish and Wildlife Service, Gainesville, FL.

Courtenay, W. R., Jr., and D. A. Hensley. 1979b. Range expansion in southern Florida of the introduced spotted tilapia, with comments on environmental impress. *Environmental Conservation* 6(2):149-151.

Courtenay, W. R., Jr., and J. R. Stauffer, Jr. 1990. The introduced fish problem and the aquarium fish industry. *Journal of the World Aquaculture Society* 21(3):145-159.

Courtenay, W. R., Jr., D. A. Hensley, J. N. Taylor, and J. A. McCann. 1986. Distribution of exotic fishes in North America. Pages 675-698 in C. H. Hocutt, and E. O. Wiley, editors. *The zoogeography of North American freshwater fishes*. John Wiley and Sons, New York, NY.

Courtenay, W. R., Jr., H. F. Sahlman, W. W. Miley, II, and D. J. Herrema. 1974. Exotic fishes in fresh and brackish waters of Florida. *Biological Conservation* 6(4):292-302.

- Gray, S.F., R.S. Smith, N.J. Reynolds, and E.W. Williams. 1990. Fish tank granuloma. *British Medical Journal* 300:1069-1070.
- Hogg, R. G. 1976. Ecology of fishes of the family Cichlidae introduced into the fresh waters of Dade County, Florida. Unpublished doctoral dissertation. University of Miami, Coral Gables, FL. 142 pp.
- Howard, B.J., J. Klaas, S.J. Rubin, A.S. Weissfeld, and R.C. Tilton. 1987. Clinical and pathogenic microbiology. C.V. Mosby, St. Louis, Missouri. 968 pp.
- Hugg, D.O. 1996. MAPFISH georeferenced mapping database. Freshwater and estuarine fishes of North America. Life Science Software. Dennis O. and Steven Hugg, 1278 Turkey Point Road, Edgewater, Maryland, USA.
- Keith, P., P.Y. Le Bail, and P. Planquette, 2000. Atlas des poissons d'eau douce de Guyane. Tome 2, Fascicule I: Batrachoidiformes, Mugiliformes, Beloniformes, Cyprinodontiformes, Synbranchiformes, Perciformes, Pleuronectiformes, Tetraodontiformes. Collection Patrimoines Naturels 43(I): 286p. Paris: Publications scientifiques du Muséum national d'Histoire naturelle.
- Kullander, S.O. 2003. Cichlidae (Cichlids). p. 605-654. In: R.E. Reis, S.O. Kullander and C.J. Ferraris, Jr. (eds.) Checklist of the Freshwater Fishes of South and Central America. Porto Alegre: EDIPUCRS, Brasil.
- Kullander, S. O. and H. Nijssen. 1989. The cichlids of Surinam, Teleostei: Labroidei. E. J. Brill, New York, NY.
- Kushlan, J. A. 1972. The exotic fish (*Aequidens portalegreensis*) in the Big Cypress Swamp. *Florida Naturalist* 45:29.
- Landsell, W., B. Dixon, N. Smith, and L. Benjamin. 1993. Isolation of several *Mycobacterium* species from fish. *Journal of Aquatic Animal Health* 5:73-76.
- Loftus, W. F. and J. A. Kushlan. 1987. Freshwater fishes of southern Florida. *Bulletin of the Florida State Museum of Biological Science* 31(4):255.
- Mills, D. and G. Vevers. 1989. The Tetra encyclopedia of freshwater tropical aquarium fishes. Tetra Press, New Jersey. 208 p.
- Nigrelli, R.F. and H. Vogel. 1963. Spontaneous tuberculosis in fishes and in other cold-blooded vertebrates with special reference to *Mycobacterium fortuitum* Cruz from fish and human lesions. *Zoologica (New York)* 48:131-144.
- Parisot, T.J. and J.N. Wood. 1970. Fish mycobacteriosis (tuberculosis). U.S. Fish and Wildlife Service Fish Disease Leaflet 7. U.S. Department of the Interior, Washington, D.C. 2 pp.

- Riehl, R. and H.A. Baensch. 1996. Aquarien Atlas, Band 1. 10th edition. Mergus Verlag GmbH, Melle, Germany. 992 p.
- Rivas, L. R. 1965. Florida fresh water fishes and conservation. Quarterly Journal of the Florida Academy of Science 28(3):255-258.
- Tilmant, J.T. 1999. Management of nonindigenous aquatic fish in the U.S. National Park System. National Park Service. 50 pp.
- USFWS. 2005. U.S. Fish and Wildlife Service, National Wildlife Refuge System, Invasive Species. Available: <http://www.nwrinvasives.com/index.asp> (Last accessed 2006).